



CHICAGO JOURNALS



---

Tutorial: Exact Solutions for the Populations of the n-level Ion

Author(s): M. Taylor and J. M. Vílchez

Reviewed work(s):

Source: *Publications of the Astronomical Society of the Pacific*, Vol. 121, No. 885 (November 2009), pp. 1257-1266

Published by: [The University of Chicago Press](#) on behalf of the [Astronomical Society of the Pacific](#)

Stable URL: <http://www.jstor.org/stable/10.1086/648121>

Accessed: 03/10/2012 07:36

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



The University of Chicago Press and Astronomical Society of the Pacific are collaborating with JSTOR to digitize, preserve and extend access to *Publications of the Astronomical Society of the Pacific*.

<http://www.jstor.org>

## Tutorial: Exact Solutions for the Populations of the $n$ -level Ion

M. TAYLOR,<sup>1</sup>

Institute for Space Applications and Remote Sensing (ISARS), National Observatory of Athens (NOA),  
Vas. Pavlou & I. Metaxa, 15236 Penteli, Greece; michael@space.noa.gr

AND

J. M. VÍLCHEZ

Departamento de Astrofísica Extragaláctica, Instituto de Astrofísica de Andalucía (IAA), CSIC,  
Camino Bajo de Huétor 50, 18008 Granada, Spain; jvm@iaa.es

Received 2009 April 29; accepted 2009 September 8; published 2009 October 20

**ABSTRACT.** This tutorial presents a review of the analytical approach to obtain exact solutions for the populations of  $n$ -level ions, and summarizes the ideas behind detailed balance and the statistical physics of collisionally-excited ions. Seaton’s analytical solution for the populations of the 3-level ion has been supplanted by matrix methods such as the master equation approach, which are now central to astronomy since there is a need to maintain a parity between improvements in quantum-mechanically calculated values for collision strengths and transition probabilities on the one hand, and three-dimensional (3D) photoionization codes used by astrophysicists for producing nebular diagnostics on the other. We show that the analytical method of solution to the problem using matrices and symbolic mathematics is straightforward, and we illustrate through theoretical, numerical, and empirical checks the validity of its results. First, we recast the equations of thermal statistical equilibrium for the energy level populations of collisionally-excited ions in the form of a well-defined matrix equation. We then show how symbolic mathematics is efficient in the inversion of equations and is able to provide the exact analytical solutions for the sought-after level populations. We present the matrices for the 5-level ion as an example of how to extend the exact solution for the 3-level ion to illustrate the general technique. We then show how the analytical results faithfully reduce to the Seaton solution when appropriate limits are taken. Spectrophotometric observations of a real ionized gas (the planetary nebula A39) are then used to obtain empirical values of forbidden line ratios and level populations for the 5-level [O III] ion. These values are compared with: (1) a best-fit 3D Monte Carlo photoionization model, and (2) the exact solution for the 5-level ion, using the symbolic mathematics approach, the exact Seaton 3-level ion solution, and a numerical approximation for the 5-level ion. It is shown that in every case, the analytical solution agrees with results obtained using observed nebular conditions to within the standard error. We provide a MATLAB code that can be adapted and tailored by astrophysicists to calculations of other  $n$ -level ions.

*Online material:* color figures

### 1. INTRODUCTION

Essentially three parameters fully determine the physical nature of an ionized nebula: the electron temperature ( $T_e$ ), the electron density ( $n_e$ ), and the level populations of the ions ( $N_i$ ). From the latter, ratios of emission rates and emission line intensities can be calculated (see, e.g., Aller 1984; Osterbrock & Ferland 2006). The 3-parameter family forms a closed set whereby knowledge of any two of them allows for a determination of the third, since

$$f(T_e, n_e, N_i) = 0. \quad (1)$$

In principle, once this equation is solved, all other secondary physical quantities such as ionization parameters, ionic abundances, and effective temperatures can then be calculated. Prior to the development of matrix solutions to the problem using the master equation method of Martin et al. (1996) or numerical approximations (Mendoza 1983), the precise functional form of  $f$  had remained elusive and unknown to astronomers and atomic physicists.

In parallel, great progress has been made in observational astrophysics, with the development of empirical tools to estimate the nebular values of  $n_e$  (Menzel et al. 1941; Copetti & Witzl, 2002; Shaw & Dufour, 1995),  $T_e$  (Shaw & Dufour, 1995; Aller et al. 1949; Seaton 1954; Spitzer 1948) and  $N_i$  (Seaton 1975). In particular, certain ratios of forbidden lines emitted by the  $p^2$ ,  $p^3$ , and  $p^4$  configuration ions present in

---

<sup>1</sup> To whom correspondence should be addressed.

the nebular gas allow the integrated electron density and electron temperature to be estimated point-to-point across projected images, while sums of forbidden lines emitted by different stages of ionization of the gas have been found to weakly correlate with ionic abundances (Shaw & Dufour, 1995; Pagel et al. 1979; Taylor & Díaz 2007).

The early work in this field was performed by Menzel, Hebb, Aller, Spitzer, Seaton, and Osterbrock (Menzel et al. 1941; Aller et al. 1949; Spitzer, 1948; Hebb & Menzel, 1940; Seaton & Osterbrock, 1957; Seaton 1960) who obtained initial estimates of the level populations of 3-level ions. An exact analytic solution for the 3-level ion was finally worked out just over 30 years ago by Seaton (1975). The computer age then allowed for the development of various higher order numerical approximations. For example, a first-order numerical approximation for the 5-level ion is currently used in nebular analysis software, such as the code FIVEL by De Robertis et al. (1987), the 3D Monte Carlo photoionization code MOCASSIN by Ercolano et al. (2003), or state of the art codes that use Breit-Pauli  $R$ -matrices to approximate ions containing even higher stages of ionization. Rodriguez (2002) has produced an approximate 34-level model of [Fe III] and Pelan & Berrington (2001) have approximated the [Fe IV] ion to 180 levels. Despite these computational achievements, only the master equation method (Martin et al. 1996) has been advanced as a way to calculate exact analytical solutions. In this tutorial, we will guide the reader through the method using a simple matrix prescription.

The theoretical context for the physics of collisionally-excited ions is the thermal equilibrium present in ionized gases. The temperature in a static nebula is fixed by the equilibrium between heating by photoionization and cooling by recombination, free-free radiation (bremsstrahlung), and line radiation. Menzel and coworkers (Menzel et al. 1941; Hebb & Menzel 1940) first set up the equation of energy conservation in ionized nebulae, whereby heating due to the energy absorbed by photoionization of the gas ( $G$ ) is balanced by cooling due to the energy liberated in capture and subsequent recombination events ( $L_R$ ), free-free Bremsstrahlung emission ( $L_{ff}$ ), and energy emitted in collisionally-excited radiative decays ( $L_C$ ),

$$G = L_R + L_{ff} + L_C. \quad (2)$$

Closed forms for  $G$ ,  $L_R$ , and  $L_{ff}$  can be found in standard texts on the physics of ionized gaseous nebula (see, e.g., Aller 1984; or Osterbrock & Ferland, 2006). The aim of this article is to show how to calculate a closed form for  $L_C$

$$L_C(N_i) = \sum_i N_i \sum_{j<i} A_{ij} h\nu_{ij}, \quad (3)$$

the cooling due to the emission of collisionally-excited lines (CELs) with level populations  $N_i$ , transition probabilities  $A_{ij}$ , de-excitation rate coefficients  $q_{ij}$ , and energy level potential differences  $h\nu_{ij}$ . In this tutorial, we will focus on the low-density regime below the critical density  $N_c(i) = \sum_{j<i} A_{ij} / \sum_{j \neq i} q_{ij}$ ,

where collisional de-excitation of level  $i$  is negligible—the thermal regime of the vast majority of planetary nebulae.

## 2. THE EXACT SOLUTION METHOD

In many gaseous nebulae, mechanical or magnetic energy from hydromagnetic waves is dissipated in the gas and there can exist regions where the mean temperature is raised such that atomic levels are excited through collisions with electrons. However, in the low density plasmas of many nebulae and ionized hydrogen (H II) regions, the most frequently observed optical lines come from the *forbidden* lines that violate the Laporte Parity Rule (Aller 1984). For these lines, we need to know the transition probabilities ( $A_{ij}$ ) and collision strengths ( $\Omega_{ij}$ ) for the excitation of metastable levels. Forbidden lines cannot be observed in the laboratory and so we have to rely entirely on theory to determine  $A_{ij}$  and  $\Omega_{ij}$  (see, e.g., Mendoza 1983). In what follows, we use the matrix master equation to rigorously solve for the populations of the  $n$ -level ion that give rise to these emitted forbidden lines.

One should start with the formal general equations of statistical equilibrium for an ion with excited levels ( $i$ ) above ground (Aller 1984, eqs. 5–32), which can be written (Appendix A)

$$\sum_{j \neq i} N_j \frac{\Omega_{ji} \epsilon_{ij}}{\omega_j} + \sum_{j > i} N_j \frac{A_{ji}}{Kx} - \sum_{j \neq i} N_i \frac{\Omega_{ji}}{\omega_i} - \sum_{j < i} N_i \frac{A_{ij}}{Kx} = 0, \quad (4)$$

and which are subject to the total ion density condition that the sum overall level populations (i.e., over all stages of ionization) is equal to the total number density ( $N$ ) of the ions [ $\text{cm}^{-3}$ ]

$$\sum_i N_i = N. \quad (5)$$

Here we have adopted the excitation potential convention  $E_i > E_j$  such that the transition  $i \rightarrow j$  corresponds to de-excitation and  $A_{ij}$  is the transition probability per second, the Seaton variable  $x = n_e / T_e^{1/2}$  (Seaton 1954) for electron temperature  $T_e$  [ $K$ ] and electron density  $n_e$  [ $\text{cm}^{-3}$ ],  $\Omega_{ji}$  is the velocity-averaged collision strength (Seaton 1968), the statistical weights are  $\omega_k = (2k + 1)$ , and the atomic constant  $K = 8.629 \times 10^{-6}$ .

The full statistical equilibrium is therefore described by a set of  $(n - 1)$  simultaneous equations in  $n$  unknowns for the excited levels above the ground level ( $i = 1$ ), with closure ensured through the addition of the total ion density condition. In matrix form,

$$\begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \cdots & \alpha_{2n} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} & \cdots & \alpha_{3n} \\ \alpha_{41} & \alpha_{42} & \alpha_{43} & \cdots & \alpha_{4n} \\ \alpha_{51} & \alpha_{52} & \alpha_{53} & \cdots & \alpha_{5n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \alpha_{n3} & \cdots & \alpha_{nn} \end{bmatrix} \begin{bmatrix} N_1 \\ N_2 \\ N_3 \\ N_4 \\ N_5 \\ \vdots \\ N_n \end{bmatrix} = \begin{bmatrix} N \\ 0 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad (6)$$

such that the elements  $\alpha_{ij}$  are equal to the coefficients of  $N_i$ . The first row with  $\alpha_{1j} = 1$  is the total ion density condition. Introducing the matrix  $\tilde{\mathbf{A}}$  of coefficients ( $\alpha_{ij}$ ), the vector  $\mathbf{y}$  of level populations ( $N_i$ ), and the vector  $\mathbf{b}$  for the righthand side, the matrix master equation can be written simply and succinctly as

$$\tilde{\mathbf{A}}\mathbf{y} = \mathbf{b}. \quad (7)$$

Provided that  $|\tilde{\mathbf{A}}| \neq 0$  (which is always the case since ions have nonnegative activation levels above ground), then we can use symbolic manipulation software such as MATLAB or Mathematica to obtain  $(\tilde{\mathbf{A}})^{-1}$  and hence the level population vector

$$\mathbf{y} = (\tilde{\mathbf{A}})^{-1}\mathbf{b}. \quad (8)$$

Since  $b_1 = N$  and  $b_{i>1} = 0$ , the sought-after level populations are identically given by

$$\begin{aligned} N_1 = y_1 &= (\tilde{\mathbf{A}}^{-1})_{11}N & N_2 = y_2 &= (\tilde{\mathbf{A}}^{-1})_{21}N \\ N_3 = y_3 &= (\tilde{\mathbf{A}}^{-1})_{31}N & N_4 = y_4 &= (\tilde{\mathbf{A}}^{-1})_{41}N \\ N_5 = y_5 &= (\tilde{\mathbf{A}}^{-1})_{51}N & \vdots & \\ N_n = y_n &= (\tilde{\mathbf{A}}^{-1})_{n1}N. \end{aligned} \quad (9)$$

It is at this point that we can identify the function  $f$  in equation (1). We know from the above definitions that the population of the  $i$ th level  $N_i$  is given by

$$N_i = (\tilde{\mathbf{A}}^{-1})_{i1}N. \quad (10)$$

If we use now introduce the matrix of cofactors  $\tilde{\mathbf{C}}$  of  $\tilde{\mathbf{A}}$ , then its transpose  $\tilde{\mathbf{C}}^T$  (the adjugate matrix) allows for an algebraic solution for the matrix inverse

$$(\tilde{\mathbf{A}}^{-1}) = \frac{\tilde{\mathbf{C}}^T}{|\tilde{\mathbf{A}}|}, \quad (11)$$

such that the level populations will have the general form

$$N_i = \frac{N}{|\tilde{\mathbf{A}}|} C_{i1}[x(n_e, T_e), \Omega_{ji}(T_e), \epsilon_{ij}(T_e)], \quad (12)$$

or equivalently,

$$N_i = f(n_e, T_e) \quad (13)$$

as implied by equation (1).

Equations (4)–(9) allow the astrophysicist to model (exactly) an ion of as many levels as required.

### 3. RESULTS

#### 3.1. The 5-Level Ion

Ions having  $p^2$ ,  $p^3$ , and  $p^4$  electron configurations all have five low-lying energy levels. For such ions, collisional and

radiative transitions can occur between any of the levels, and excitation and de-excitation cross sections as well as collision strengths exist between all pairs of levels. A central assumption that is often made is that line emission from these five levels alone provides a good approximation to the expected emission lines of the full  $n$ -level ion. The justification (see, e.g., Osterbrock & Ferland 2006) is that higher levels in these ions are not significantly populated through collisions, recombinations, or other mechanisms. Although this is true for weakly ionized, low-density nebular plasmas such as planetary nebulae and HII regions, active galactic nuclei (AGNs) and supernovae (SNs) are much more strongly ionized, meaning that higher level transitions are likely to be significant to their emission line spectra—as evidenced by the tremendous numerical modeling work of the IRON and OPACITY projects.

For a 5-level ion in a steady state with  $E_5 > E_4 > E_3 > E_2 > E_1$ , the total number density condition for each ion species is given by  $N_1 + N_2 + N_3 + N_4 + N_5 = N$  and the equations of statistical equilibrium (4) give rise to the following four exact level population equations for the excited levels above ground

$$\begin{aligned} N_1 \frac{\Omega_{21}\epsilon_{21}}{3} - N_2 \left( \frac{A_{21}}{Kx} + \frac{\Omega_{12} + \Omega_{32} + \Omega_{42} + \Omega_{52}}{5} \right) \\ + N_3 \left( \frac{A_{32}}{Kx} + \frac{\Omega_{32}\epsilon_{23}}{7} \right) + N_4 \left( \frac{A_{42}}{Kx} + \frac{\Omega_{42}\epsilon_{24}}{9} \right) \\ + N_5 \left( \frac{A_{52}}{Kx} + \frac{\Omega_{52}\epsilon_{25}}{11} \right) = 0 \\ N_1 \frac{\Omega_{13}\epsilon_{31}}{3} + N_2 \frac{\Omega_{23}\epsilon_{32}}{5} \\ - N_3 \left( \frac{A_{31} + A_{32}}{Kx} + \frac{\Omega_{13} + \Omega_{23} + \Omega_{43} + \Omega_{53}}{7} \right) \\ + N_4 \left( \frac{A_{43}}{Kx} + \frac{\Omega_{43}\epsilon_{34}}{9} \right) + N_5 \left( \frac{A_{53}}{Kx} + \frac{\Omega_{53}\epsilon_{35}}{11} \right) = 0 \\ N_1 \frac{\Omega_{14}\epsilon_{41}}{3} + N_2 \frac{\Omega_{24}\epsilon_{42}}{5} + N_3 \frac{\Omega_{34}\epsilon_{43}}{7} \\ - N_4 \left( \frac{A_{41} + A_{42} + A_{43}}{Kx} \right) \\ - N_4 \left( \frac{\Omega_{14} + \Omega_{24} + \Omega_{34} + \Omega_{54}}{9} \right) \\ + N_5 \left( \frac{A_{54}}{Kx} + \frac{\Omega_{54}\epsilon_{45}}{11} \right) = 0 \\ N_1 \frac{\Omega_{15}\epsilon_{51}}{3} + N_2 \frac{\Omega_{25}\epsilon_{52}}{5} + N_3 \frac{\Omega_{35}\epsilon_{53}}{7} + N_4 \frac{\Omega_{45}\epsilon_{54}}{9} \\ - N_5 \left( \frac{A_{51} + A_{52} + A_{53} + A_{54}}{Kx} \right) \\ - N_5 \left( \frac{\Omega_{15} + \Omega_{25} + \Omega_{35} + \Omega_{45}}{11} \right) = 0. \end{aligned} \quad (14)$$

The elements  $\alpha_{ij}$  of  $\tilde{\mathbf{A}}$  are then equal to the coefficients of the level populations  $N_i$  and are listed in Appendix B. We have deliberately not inserted the values of  $A_{ji}$ ,  $\Omega_{ji}$ , and  $\epsilon_{ij}$  up until now as they are estimated values that will evolve with the accuracy of quantum mechanical calculations currently being performed by the IRON and OPACITY projects.

The matrix equation  $\tilde{\mathbf{A}}\mathbf{y} = \mathbf{b}$  having the coefficients  $\alpha_{ij}$  is exact for the 5-level ion. At this stage, the problem is reduced to finding the matrix inverse. This can be done numerically as in Martin et al. (1996) or using singular value decomposition. With the advent of symbolic mathematics software such as MATLAB and MAPLE, it is also possible to find the inverse matrix  $\tilde{\mathbf{A}}^{-1}$  analytically so as to obtain exact expressions for the level populations. For example, the full solution includes terms up to  $(Kx)^4$ , and while cumbersome and too large to reproduce here, is exact. However, in Appendix C we provide a general public license (GPL) MATLAB code for generation of the symbolically-obtained analytic solutions for comparison with exact numerical solutions. In general, an  $n$ -level ion will contain terms up to  $(Kx)^{n-1}$ . The inverse matrix solution for the 3-level ion includes terms only up to  $(Kx)^2$  and we show in § 3.2 how the known analytic solution (Seaton 1975) can be reproduced.

### 3.2. The 3-Level Ion

For ions of type  $p^2$  such as the [O III] ion shown in Figure 1, suppression of all terms with indices equal to 4 and 5 reduces the general solution for the 5-level ion to that of the 3-level ion. This amounts to removing the fine splitting physics of the  $^3P$  level. As a consequence, the excitation potentials  $E_2$  and  $E_3$  of the 3-level ion correspond identically with  $E_4$  and  $E_5$  of the 5-level ion while  $E_1 = 0$  eV is the ground level. Similarly, a suitable suppression of indices allows the 5-level representations of the  $p^3$  and  $p^4$  ions to be reduced to their exact 3-level counterparts. In every case, the following exact level populations for the 3-level ion are obtained

$$N_1 = \frac{3N}{D} \left[ 35A_{21}A_{31} + 35A_{21}A_{32} + Kx \left( 5A_{21}\Omega_{13} + 5A_{21}\Omega_{23} + 7A_{31}\Omega_{12} + 7A_{31}\Omega_{32} + 7A_{32}\Omega_{12} + 7A_{32}\Omega_{32} - 7A_{32}\Omega_{23}\epsilon_{32} \right) + K^2x^2 \left( \Omega_{12}\Omega_{13} + \Omega_{12}\Omega_{23} + \Omega_{13}\Omega_{32} + \Omega_{23}\Omega_{32} - \Omega_{23}\epsilon_{32}\Omega_{32}\epsilon_{23} \right) \right] \quad (15)$$

$$N_2 = \frac{5N}{D} \left[ 7Kx \left( A_{31}\Omega_{12}\epsilon_{21} + A_{32}\Omega_{12}\epsilon_{21} + A_{32}\Omega_{13}\epsilon_{31} \right) + K^2x^2 \left( \Omega_{12}\epsilon_{21}\Omega_{13} + \Omega_{12}\epsilon_{21}\Omega_{23} + \Omega_{13}\epsilon_{31}\Omega_{32}\epsilon_{23} \right) \right] \quad (16)$$

$$N_3 = \frac{7N}{D} \left[ Kx \left( 5A_{21}\Omega_{13}\epsilon_{31} \right) + K^2x^2 \left( \Omega_{12}\epsilon_{21}\Omega_{23}\epsilon_{32} + \Omega_{12}\Omega_{13}\epsilon_{31} + \Omega_{13}\Omega_{31}\Omega_{32} \right) \right] \quad (17)$$

with denominator  $D$ ,

$$D = 105A_{21}A_{31} + 105A_{21}A_{32} + Kx \left[ 15A_{21}\Omega_{13} + 15A_{21}\Omega_{23} + 21A_{31}\Omega_{12} + 21A_{31}\Omega_{32} + 21A_{32}\Omega_{12} + 21A_{32}\Omega_{32} - 21A_{32}\Omega_{23}\epsilon_{32} + 35A_{21}\Omega_{13}\epsilon_{31} + 35A_{31}\Omega_{12}\epsilon_{21} + 35A_{32}\Omega_{12}\epsilon_{21} + 35A_{32}\Omega_{13}\epsilon_{31} \right] + K^2x^2 \left[ 3\Omega_{12}\Omega_{13} + 3\Omega_{12}\Omega_{23} + 3\Omega_{13}\Omega_{32} + 3\Omega_{23}\Omega_{32} - 3\Omega_{23}\epsilon_{32}\Omega_{32}\epsilon_{23} + 5\Omega_{12}\epsilon_{21}\Omega_{13} + 5\Omega_{12}\epsilon_{21}\Omega_{23} + 5\Omega_{13}\epsilon_{31}\Omega_{32}\epsilon_{23} + 7\Omega_{12}\Omega_{13}\epsilon_{31} + 7\Omega_{12}\epsilon_{21}\Omega_{23}\epsilon_{32} + 7\Omega_{13}\epsilon_{31}\Omega_{32} \right]. \quad (18)$$

In order to compare the exact matrix master equation solution for the  $n = 3$  level ion with the known analytic solution of Seaton (1975), we consider the ratio of the level populations  $N_3$  and  $N_2$  (although nowadays the inverse  $N_2/N_3$  for the ratio of nebular to auroral lines is customary)

$$N_3/N_2 = [7N/D][Kx(5A_{21}\Omega_{13}\epsilon_{31}) + K^2x^2(\Omega_{12}\epsilon_{21}\Omega_{23}\epsilon_{32} + \Omega_{12}\Omega_{13}\epsilon_{31} + \Omega_{13}\Omega_{31}\Omega_{32})]/[5N/D][7Kx(A_{31}\Omega_{12}\epsilon_{21} + A_{32}\Omega_{12}\epsilon_{21} + A_{32}\Omega_{13}\epsilon_{31}) + K^2x^2(\Omega_{12}\epsilon_{21}\Omega_{13} + \Omega_{12}\epsilon_{21}\Omega_{23} + \Omega_{13}\epsilon_{31}\Omega_{32}\epsilon_{23})].$$

Extracting the term  $(\Omega_{13}\epsilon_{31})$  from the numerator and  $(\Omega_{12}\epsilon_{21})$  from the denominator and canceling like terms we obtain

$$\frac{N_3}{N_2} = \frac{\Omega_{13}\epsilon_{31}A_{21}}{\Omega_{12}\epsilon_{21}A_{32}} C \frac{[1 + \frac{Kx}{A_{21}}\Psi_3]}{[1 + \frac{Kx}{A_{32}}\Psi_2]}, \quad (19)$$




FIG. 1.—Five-level [O III] diagram. Radiative transitions marked by arrows have their central emission wavelength in Å. See the electronic edition of the PASP for a color version of this figure.



where

$$C = \frac{1}{1 + \frac{A_{31}}{A_{32}} + \frac{\Omega_{13}\epsilon_{31}}{\Omega_{12}\epsilon_{21}}}$$

$$\Psi_2 = C \left( \frac{\Omega_{12}\epsilon_{21}\Omega_{13} + \Omega_{12}\epsilon_{21}\Omega_{23} + \Omega_{13}\epsilon_{31}\Omega_{32}\epsilon_{23}}{7\Omega_{12}\epsilon_{21}} \right) \quad (20)$$

$$\Psi_3 = \frac{\Omega_{12}\epsilon_{21}\Omega_{23}\epsilon_{32} + \Omega_{12}\Omega_{13}\epsilon_{31} + \Omega_{13}\epsilon_{31}\Omega_{32}\epsilon_{23}}{5\Omega_{13}\epsilon_{31}}.$$

This is the same result as that obtained by Seaton for the exact 3-level ion (Seaton 1975, eq. 1.11).

By considering the dimensions of the physical variables (there are 7 metric units: mass  $M$ [Kg], length  $L$ [m], time  $T$ [s], temperature  $\theta$ [K], electrical current  $I$ [A], concentration  $N$ [mol], and light intensity  $J$ [cd]), a further theoretical check is provided by the requirement for dimensional homogeneity which for a set of physical variables  $\{x_i\}$  requires that  $[x_1^{\epsilon_1} x_2^{\epsilon_2} \dots x_n^{\epsilon_n}] = 1$  in order to satisfy causality (Taylor et al. 2008). The dimensions of the various physical quantities and parameters presented in our derivation are

$$\begin{aligned} [n_e] &= [N_i] = L^{-3} \\ [q_{ji}] &= [q_{ij}] = L^3 T^{-1} \\ [n_e N_i q_{ji}] &= L^{-3} T^{-1} \\ [A_{ji}] &= T^{-1} \\ [x(n_e, T_e)] &= L^{-3} \Theta^{-1/2} \\ [K] &= L^3 T^{-1} \Theta^{1/2} \\ [\epsilon_{ji}] &= [\omega_j] = [\Omega_{ji}] = [\alpha_{ij}] = 1, \end{aligned}$$

such that

$$\left[ \frac{A_{ji}}{Kx} \right] = \left[ \frac{Kx}{A_{ji}} \right] = [C] = [\Psi_2] = [\Psi_3] = 1.$$

All equations obtained from the matrix master equation approach were thoroughly checked for dimensional homogeneity with these dimensional relations and no inconsistencies were found.

### 3.3. Empirical Checks

In order to compare the algebraically-obtained for the level populations of the 5-level ion with those obtained from observational measurements and numerical computer simulations, it is necessary to know the values of the measured emission line intensities of an ion  $I(i, k)$  and the average nebular electron density  $n_e$  and electron temperature  $T_e$ , as well as the quantum-mechanically calculated values of  $A_{ji}$ ,  $\epsilon_{ij}(T_e)$ ,  $\Omega_{ji}(T_e)$ , and  $x(n_e, T_e)$ . The line intensities (being the only direct observable) are proportional to the line emission rate  $j(i, k)$  of line photons resulting from downward transitions  $i \rightarrow k$ ,

$$4\pi j(i, k) = A_{ik} N(X^l) N_i h\nu_{ik} \quad (21)$$

where  $X^l$  is the ion species; e.g.,  $O^{+2}$  and  $\Delta E = h\nu_{ik}$  corresponds to the energy difference between excitation potentials  $E_i$  and  $E_k$ . Hence, a ratio of line intensities will involve only the ratio of level populations  $N_i$  and ionic constants  $A_{ik}$  and  $\Delta E_{ik}$ . So, for example, the ratio of auroral to nebular lines of [O III] which we denote  $R[O III]$

$$R[O III] \equiv \frac{I(\lambda 4959) + I(\lambda 5007)}{I(\lambda 4363)}, \quad (22)$$

for the 3-level ion is given by

$$\begin{aligned} R[O III] &= \frac{j(2, 1)_{5007} + j(2, 1)_{4959}}{j(3, 2)_{4363}} \\ &= \frac{N_2 \Delta E_{21}}{N_3 \Delta E_{32}} \left[ \frac{A_{21}(5007) + A_{21}(4959)}{A_{32}(4363)} \right], \quad (23) \end{aligned}$$

and for the 5-level ion is

$$R[O III] = \frac{N_4}{N_5} \left[ \frac{\Delta E_{43} A_{43} + \Delta E_{42} A_{42}}{\Delta E_{54} A_{54}} \right]. \quad (24)$$

Therefore, by measuring  $R[O III]$  observationally, it is possible to compare its value with that obtained using the level populations (9) provided by the matrix master equation method using the values of  $A_{ji}$ ,  $\epsilon_{ij}(T_e)$ ,  $\Omega_{ji}(T_e)$ , and  $x(n_e, T_e)$  evaluated at the measured value of electron density and temperature.

### 3.4. Spectrophotometry of the Spherical Planetary Nebula A39

The spherical planetary nebula A39 was discovered by George Abell in his 1957 survey of the Southern Hemisphere (Abell 1966) and has recently been subject to thorough spectrophotometry (Jacoby et al. 2000). From these detailed measurements made by George Jacoby and coworkers at Kitt Peak, the reddening-corrected line ratio  $R[O III]$  averaged over the whole nebula was measured to be (Jacoby et al. 2000)

$$R[O III] = \frac{3.98 \pm 0.182 + 11.31 \pm 0.524}{0.24 \pm 0.019} = 64.24 \pm 6.58 \quad (25)$$

with all line intensities  $I(\lambda)$  relative to  $I(H_\beta)$ . The electron density and electron temperature were estimated to be  $n_e \approx 30 \text{ cm}^{-3}$   $T_e \approx 15400 \text{ K}$  (Jacoby et al. 2000). Inserting the most up-to-date values of the atomic and ionic constants  $A_{ij}$ ,  $\Omega_{i,j}$ , and  $\epsilon_{ji}$  (see Osterbrock & Ferland 2006 and references therein) calculated at  $n_e = 30 \text{ cm}^{-3}$  and  $T_e = 15400 \text{ K}$ , we obtain  $R[O III] = 65.47$  for the exact 5-level [O III] ion and  $R[O III] = 64.92$  for the Seaton 3-level [O III] ion. Furthermore, the first-order numerical approximation to the 5-level [O III] ion provided by the code TEMDEN (De Robertis et al. 1987) gives

$R[\text{O III}] = 65.16$  at this electron density and temperature; all well within the observational standard error.

### 3.5. Three-Dimensional Photoionization Modeling of A39

Using the 3D Monte Carlo photoionization code MOCASSIN (Ercolano et al. 2003) and a best-fit model to the observed spectrum with a 3-shell density profile (Figure 2), one obtains the estimate  $R[\text{O III}] = 66.57$  at an average nebular electron density  $\langle n_e \rangle = 11.94 \text{ cm}^{-3}$  and temperature  $\langle T_e \rangle = 15896 \text{ K}$ . Once again, inserting the values of the atomic and ionic constants  $A_{ij}$ ,  $\Omega_{ij}$ , and  $\epsilon_{ji}$  calculated at  $n_e = 11.94 \text{ cm}^{-3}$  and  $T_e = 15896 \text{ K}$ , we obtained the values  $R[\text{O III}] = 61.98$  for the exact 5-level [O III] ion,  $R[\text{O III}] = 61.02$  for the Seaton 3-level [O III] ion, and  $R[\text{O III}] = 60.93$  with TEMDEN. Incidentally, a rerun of MOCASSIN using the value of averaged electron density  $\langle n_e \rangle = 30 \text{ cm}^{-3}$ , suggested from the observational study of Jacoby and coworkers, yielded a slightly lower average nebular temperature  $\langle T_e \rangle = 14997 \text{ K}$ . In this particular case, MOCASSIN obtained the much higher value  $R[\text{O III}] = 69.91$ , in agreement also with high values for the exact 5-level ion  $R[\text{O III}] = 70.25$ , the Seaton 3-level ion  $R[\text{O III}] = 69.22$ , and TEMDEN  $R[\text{O III}] = 69.05$ . To check that such an increase is to be expected at lower electron temperatures, we calculated the value of  $R[\text{O III}]$  using the tabulated values for [O III] (Lennon & Burke 1994) of  $A_{ij}$ ,  $\Omega_{ij}$ , and  $\epsilon_{ji}$  at an electron density  $n_e = 30 \text{ cm}^{-3}$  and temperature  $T_e = 10000 \text{ K}$ . Here, we found that the exact 5-level ion gives  $R[\text{O III}] = 213.40$ , the Seaton 3-level ion gives  $R[\text{O III}] = 203.11$ , and the approximate 5-level ion calculation of TEMDEN gives  $R[\text{O III}] = 209.77$ ; all dramatically larger and much more dispersed. The level populations, as expected, are very sensitive to electron temperature. All of these results are collected and presented in Table 1 for ease of comparison.

At electron densities of  $n_e = 30 \text{ cm}^{-3}$ , all of the results are within the observational standard error apart from the low-temperature case ( $T_e = 10000$ ) where, even here, there is consistency between the exact 5-level ion, the Seaton 3-level ion, and the numerical approximation to the 5-level ion provided by TEMDEN. In the case of the best-fit 3D photoionization model with MOCASSIN using a 3-shell density profile having an average electron density of  $\langle n_e \rangle = 11.94 \text{ cm}^{-3}$ , the results are more dispersed with the code producing a value of  $R[\text{O III}]$  closer to the observations. This may be due to uncertainties in the estimated values of the atomic constants  $A_{ij}$ ,  $\Omega_{ij}$ , and  $\epsilon_{ji}$ . What is interesting is that the value of  $R[\text{O III}]$  from the exact 5-level ion is consistently higher than both the exact Seaton 3-level ion and the numerical approximation to the 5-level ion of TEMDEN, suggesting that the extra terms are not negligible, particularly at lower electron temperatures.

## 4. DISCUSSION

This tutorial presents the master equation method for the calculation of the populations of the  $n$ -level ion from the general equations of statistical equilibrium. The matrix formulation presented here, together with the algebraic power offered by symbolic manipulation software, means that it is possible to obtain an exact solution for the populations of ions of an arbitrary number of levels undergoing collisional excitation and de-excitation for comparison with other exact numerical inversions. This is particularly important in studies of strongly ionized ions in high-energy objects such as AGNs or SNs.

The three-fold comparison of theory with observations, simulations, and numerical approximations gives consistent results and we invite the reader to extend the application to other ions (besides [O III] considered here). The setting up of the  $n$ -level problem is straightforward and we hope that these results,




FIG. 2.—(a) Spectrometric image of the spherical planetary nebula A39 using an [O III] 5007 filter (Jacoby et al. 2000); (b) a 3D IDL plot of the  $R[\text{O III}]$  distribution for the same nebula at an initial electron temperature  $T_e = 15400 \text{ K}$  and with a 3-shell electron density profile  $n_e$  calculated with the photoionization code MOCASSIN. See the electronic edition of the *PASP* for a color version of this figure.

TABLE 1  
 VALUES OF THE [O III] LINE RATIO  $R[O III]$  OBTAINED FROM THE SPHERICAL PLANETARY NEBULAE A39

Case	$n_e$	$T_e$	$R[O III]$				
			Observed	SIM	5-level	3-level	NUM
I	$n_e \approx 30$	$T_e \approx 15400$	$64.24 \pm 6.58$		65.47	64.92	65.16
II	$\langle n_e \rangle = 11.94$	$\langle T_e \rangle = 15896$		66.57	61.98	61.02	60.93
	$\langle n_e \rangle = 30$	$\langle T_e \rangle = 14997$		69.91	70.25	69.22	69.05
III	$n_e = 30$	$T_e = 10000$			213.40	203.11	209.77

NOTE.—Observed, simulated (SIM), exact 5-level, and analytical Seaton 3-level solutions, and a first-order numerical approximation (NUM) to the 5-level ion using the code TEMDEN. Case studies: I, Spectrophotometry of A39; II, 3D MOCASSIN photoionization models of A39; III, quantum-mechanical calculations of  $\Omega_{ji}$ .

in conjunction with the supplementary MATLAB code, will help make the incorporation of higher order transitions into existing astrophysics software easier for the study of very high-ionization objects such as AGNs and SNs. In addition, the interested reader may also like to take up the challenge of including more complex atomic processes such as fluorescence and recombination mechanisms.

M. T. would like to thank Luis Colina Robledo for kindly making available a copy of Aller's 1984 book, *Physics of*

*Thermal Gaseous Nebulae*, José Cernicharo for originally facilitating this work, the Greek National Scholarships Foundation (IKY) for their financial support, and to the members of DAMIR-CSIC (Madrid) and ISARS-NOA (Athens) for their hospitality. This work is partially funded by project AYA2007-67965-C03-02 of the Spanish MCINN. This work is dedicated to the memory of Donald E. Osterbrock, Bernard E. J. Pagel, and Michael J. Seaton who have done so much to advance this important field.

## APPENDIX A

### A REFORMULATION OF THE GENERAL EQUATIONS OF STATISTICAL EQUILIBRIUM

Assuming that level populations  $N_i$  are determined by spontaneous emission (with transition probability  $A_{ij} s^{-1}$ ), then the formal general equations of statistical equilibrium for an ion with excited levels ( $i$ ) above ground having excitation rate coefficients ( $q_{ji}$ ) and de-excitation rate coefficients ( $q_{ij}$ ) to the other levels  $j \neq i$  are given by Menzel et al. (1941: eqs. 5–32)

$$\sum_{j \neq i} N_j n_e q_{ji} + \sum_{j > i} N_j A_{ji} = \sum_{j \neq i} N_i n_e q_{ij} + \sum_{j < i} N_i A_{ij},$$

subject to the total ion density condition that the sum over all level populations (i.e., over all stages of ionization) is equal to the total number density ( $N$ ) of the ions [ $\text{cm}^{-3}$ ]

$$\sum_i N_i = N. \quad (\text{A1})$$

Here we have adopted the excitation potential convention  $E_i > E_j$  such that the transition  $i \rightarrow j$  corresponds to de-excitation. The next step is to note that  $q_{ij}$  times the electron density is equal to the collisional excitation rate  $c_{ij}$  [ $s^{-1}$ ]

$$n_e q_{ij} = c_{ij} \equiv Kx \frac{\Omega_{ji}}{\omega_i}, \quad (\text{A2})$$

where  $K = 8.629 \times 10^{-6}$ , the Seaton variable  $x = n_e/T_e^{1/2}$  (Copetti & Writzl 2002) for electron temperature  $T_e$  [K] and

electron density  $n_e$  [ $\text{cm}^{-3}$ ],  $\omega_i = (2i + 1)$  are the statistical weights, and where  $\Omega_{ji}$  is the velocity-averaged collision strength (Shaw & Dufour 1995)

$$\Omega_{ji} = \int_0^\infty \Omega(ji; E) e^{-E/k_B T_e} d\left(\frac{E}{k_B T_e}\right), \quad (\text{A3})$$

for colliding electrons having initial kinetic energy  $E = mu^2/2$ . The collision strengths must be calculated quantum mechanically, and consist, in general, of a part that varies slowly with energy superimposed with resonance contributions that vary rapidly. The fact that integration over a broad Maxwellian distribution of electron energies tends to dramatically smooth out the variations means that  $\Omega_{ji}$  is now known to be fairly insensitive to temperature (Aller et al. 1949). Similarly, the collisional deexcitation rate

$$c_{ji} = n_e q_{ji}, \quad (\text{A4})$$

is related to the collisional excitation rate through

$$c_{ji} = c_{ij} \frac{\omega_i}{\omega_j} \epsilon_{ij}, \quad (\text{A5})$$

with  $\epsilon_{ij} = e^{-(E_i - E_j)/k_B T_e}$ , and where  $E_i$  and  $E_j$  are the excitation potentials of the relevant levels. The radiative transition



probabilities  $A_{ij}$  are independent constants being inversely proportional to the occupancy lifetimes (and hence the energy) of the upper levels. Replacing all  $n_e q_{ji}$  and  $n_e q_{ij}$  terms in the equations of thermal equilibrium for the excited levels by equations (A1)–(A5) for the collisional excitation and de-excitation

rates, and dividing through by  $Kx (\neq 0)$ , we obtain the equation of statistical equilibrium (4) used in the main text,

$$\sum_{j \neq i} N_j \frac{\Omega_{ji} \epsilon_{ij}}{\omega_j} + \sum_{j > i} N_j \frac{A_{ji}}{Kx} - \sum_{j \neq i} N_i \frac{\Omega_{ji}}{\omega_i} - \sum_{j < i} N_i \frac{A_{ij}}{Kx} = 0.$$

## APPENDIX B

### THE MATRIX COEFFICIENTS FOR THE 5-LEVEL ION

From the four level population equations for the excited levels above ground (eq. [14]), the elements  $\alpha_{ij}$  of  $\tilde{\mathbf{A}}$  are equal to the coefficients of the level populations  $N_i$ . For the 5-level ion these are:

$$\begin{aligned} \alpha_{11} &= 1 & \alpha_{12} &= 1 & \alpha_{13} &= 1 \\ \alpha_{14} &= 1 & \alpha_{15} &= 1 & & \\ \alpha_{21} &= \frac{\Omega_{12} \epsilon_{21}}{3} & \alpha_{22} &= -\frac{A_{21}}{Kx} - \frac{\Omega_{12} + \Omega_{32} + \Omega_{42} + \Omega_{52}}{5} \\ \alpha_{23} &= \frac{A_{32}}{Kx} + \frac{\Omega_{32} \epsilon_{23}}{7} & \alpha_{24} &= \frac{A_{42}}{Kx} + \frac{\Omega_{42} \epsilon_{24}}{9} \\ \alpha_{25} &= \frac{A_{52}}{Kx} + \frac{\Omega_{52} \epsilon_{25}}{11} & \alpha_{31} &= \frac{\Omega_{13} \epsilon_{31}}{3} \\ \alpha_{32} &= \frac{\Omega_{23} \epsilon_{32}}{5} \end{aligned}$$

$$\begin{aligned} \alpha_{33} &= -\frac{(A_{31} + A_{32})}{Kx} - \frac{\Omega_{13} + \Omega_{23} + \Omega_{43} + \Omega_{53}}{7} \\ \alpha_{34} &= \frac{A_{43}}{Kx} + \frac{\Omega_{43} \epsilon_{34}}{9} & \alpha_{35} &= \frac{A_{53}}{Kx} + \frac{\Omega_{53} \epsilon_{35}}{11} \\ \alpha_{41} &= \frac{\Omega_{14} \epsilon_{41}}{3} & \alpha_{42} &= \frac{\Omega_{24} \epsilon_{42}}{5} & \alpha_{43} &= \frac{\Omega_{34} \epsilon_{43}}{7} \\ \alpha_{44} &= -\frac{(A_{41} + A_{42} + A_{43})}{Kx} - \frac{\Omega_{14} + \Omega_{24} + \Omega_{34} + \Omega_{54}}{9} \\ \alpha_{45} &= \frac{A_{54}}{Kx} + \frac{\Omega_{54} \epsilon_{45}}{11} & \alpha_{51} &= \frac{\Omega_{15} \epsilon_{51}}{3} & \alpha_{52} &= \frac{\Omega_{25} \epsilon_{52}}{5} \\ \alpha_{53} &= \frac{\Omega_{35} \epsilon_{53}}{7} & \alpha_{54} &= \frac{\Omega_{45} \epsilon_{54}}{9} \\ \alpha_{55} &= \frac{(A_{51} + A_{52} + A_{53} + A_{54})}{Kx} \\ &+ \frac{\Omega_{15} + \Omega_{25} + \Omega_{35} + \Omega_{45}}{11}. \end{aligned} \quad (\text{B1})$$

## APPENDIX C

### A MATLAB CODE FOR CALCULATING THE EXACT SOLUTIONS FOR THE $n$ -LEVEL ION

Here, we provide a general public license (GPL) MATLAB code for the generation of the matrix master equation solution of  $n$ -level ions using symbolic mathematics inversions. The code includes relevant atomic data for [O III] and, using the prescription given by equations (4)–(9), can be easily adapted for an ion of  $n$ -levels or for other ion species.

```
% PROGRAM: n_level_ion.m
%
% m-file containing the symbolic
% computations for calculation
% of the populations of the n-level ion
% with full matrix components
% provided for the exact 5-level ion and
% the Seaton 3-level ion.
%
% Written by Michael Taylor 09 October
% 2007
%
```

```
% Code distributed under the General
% Public License (GPL)

syms O12 O13 O23 E23;
syms O21 O31 O41 O51 O32 O42 O52 O43 O53
O54;
syms A21 A31 A41 A51 A32 A42 A52 A43 A53
A54;
syms E21 E31 E41 E51 E32 E42 E52 E43 E53
E54;
syms K x;
syms A y b I N;
syms Level N_total;

% Physical constants (erg, Angstrom)
ang=1e-10;
erg=1e-07;
ev=1.60217733e-19;
c=2.99792458e+08/ang;
```

```

h=6.62607554e-34/erg;
K_b=1.380658e-23;

% 5-level [OIII]: level energies above
ground (eV) (NIST)
L1=0.0;
L2=0.0140323;
L3=0.0379607;
L4=2.513565;
L5=5.354349;
L21=L2-L1;
L31=L3-L1;
L41=L4-L1;
L51=L5-L1;
L32=L3-L2;
L42=L4-L2;
L52=L5-L2;
L43=L4-L3;
L53=L5-L3;
L54=L5-L4;

% 3_level ion: Exact solution by symbolic
manipulation
% b=[N,0,0]';
% A=[1,1,1;
% O12*E21/3,-A21/(K*x)-(O12+O32)/5,A32/
(K*x)+O32*E23/7;
% O13*E31/3,O23*E32/5,-(A31+A32)/
(K*x)-(O13+O23)/7];
% Level=inv(A)*b;
% N_total=Level(1)+Level(2)+Level(3);
% simple(Level);
% pretty(ans);

% 5_level ion: Exact solution by symbolic
manipulation
b=[N,0,0,0,0]';
A=[1,1,1,1,1 ; O21*E21/3,-A21/(K*x)-(O21
+O32+O42+O52)/5,A32/(K*x)+O32*E32/7,A42/
(K*x)+O42*E42/9,A52/(K*x)+O52*E52/11;
O31*E31/3,O32*E32/5,-(A31+A32)/(K*x)-
(O31+O32+O43+O53)/7,A43/(K*x)+O43*E43/9,
A53/(K*x)+O53*E53/11; O41*E41/3,O42*E42/5,
O43*E43/7,-(A41+A42+A43)/(K*x)-(O41+O42
+O43+O54)/9, A54/(K*x)+O54*E54/11;
O51*E51/3,O52*E52/5,O53*E53/7,O54*E54/9,-
(A51+A52+A53+A54)/(K*x)-(O51+O52+O53
+O54)/11];
Level=inv(A)*b;
N_total=Level(1)+Level(2)+Level(3)+Level
(4)+Level(5);
simple(Level);
pretty(ans);

% Physical conditions for Table 1
ne=30.0;
% ne=11.94;
Te=15400;
% Te=14997;
% Te=15896;
% Te=10000;
% Te=15848;
N=1;
x=ne/(Te^(.5));
K=8.629e-06;
E21=exp(-L21/(K_b*Te));
E31=exp(-L31/(K_b*Te));E41=exp(-L41/
(K_b*Te));
E51=exp(-L51/(K_b*Te));
E32=exp(-L32/(K_b*Te));
E42=exp(-L42/(K_b*Te));
E52=exp(-L52/(K_b*Te));
E43=exp(-L43/(K_b*Te));
E53=exp(-L53/(K_b*Te));
E54=exp(-L54/(K_b*Te));

% Einstein rate coefficients (NIST)
A21=2.61e-05;
A31=3.17e-11;
A41=1.690e-06;
A51=0.0;
A32=9.76e-05;
A42=6.995e-03;
A52=2.268e-01;
A43=2.041e-02;
A53=6.091e-04;
A54=1.561;

% Collision cross-sections': numerical
approximation formulae (TEM DEN)
t4=Te*1.0e-4;
C1=1.835+t4*(0.3981-t4*0.06);
C2=0.2127+t4*(0.0767-0.013*t4);
O21=0.4825+t4*(0.0806-t4*0.022);
O31=0.2397+t4*(0.0381-t4*0.007);
O41=C1/9; O51=C2/9;
O32=1.1325+t4*(0.203-t4*0.05);
O42=C1/3;
O52=C2/3;
O43=C1*5/9;
O53=C2*5/9;
O54=0.3763+t4*(0.3375-t4*0.105);

% 5-level ion calculation
n1=subs(Level(1));
n2=subs(Level(2));

```

```
n3=subs (Level (3) );
n4=subs (Level (4) );
n5=subs (Level (5) );
```

```
R_OIII=(n4*A42*L42+n4*A43*L43) /
(n5*A54*L54);
% END: n_level_ion.m
```

## REFERENCES

- Abell, G. O. 1966, *AJ*, 144, 259
- Aller, L. H. 1984, *Physics of Thermal Gaseous Nebulae* (Dordrecht: Reidel)
- Aller, L. H., Ufford, C. W., & Van Vleck, J. H. 1949, *AJ*, 109, 42
- Copetti, M. V. F., & Writzl, B. C. 2002, *A&A*, 382, 282
- De Robertis, M. M., Dufour, R. J., & Hunt, R. W. 1987, *JRASC*, 81, 195
- Ercolano, B., Barlow, M. J., Storey, P. J., & Liu, X. W. 2003, *MNRAS*, 340, 1136
- Hebb, M. H., & Menzel, D. H. 1940, *AJ*, 92, 408
- Jacoby, G., Ferland, G. J., & Korista, K. T. 2000, *A&AS*, 197, 616
- Lennon, D. J., & Burke, V. M. 1994, *A&AS*, 103, 273
- Martin, P. G., Schwarz, D. H., & Mandy, M. E. 1996, *AJ*, 461, 265
- Mendoza, C. 1983, *IAU Symp. Proc.* 103, *Planetary Nebulae*, (Dordrecht: Reidel) 143
- Menzel, D. H., Aller, L. H., & Hebb, M. H. 1941, *AJ*, 93, 230
- Osterbrock, D. E., & Ferland, G. J. 2006, *Mercury*, 35, 40, University Science Books
- Pagel, B. E. J., Edmunds, M. G., Blackwell, D. E., Chun, M. S., & Smith, G. 1979, *MNRAS*, 189, 95
- Pelan, J. C., & Berrington, K. A. 2001, *A&A*, 365, 258
- Rodriguez, M. 2002, *A&A*, 389, 556
- Seaton, M. J. 1954, *Ann. d'Astrophys.*, 17, 74
- . 1975, *MNRAS*, 170, 475
- . 1960, *Rep. Prog. Phys.*, 23, 313
- . 1968, *Adv. AMP*, 4, 331
- Seaton, M. J., & Osterbrock, D. E. 1957, *AJ*, 125, 66
- Shaw, R. A., & Dufour, R. J. 1995, *PASP*, 107, 896
- Spitzer, L. 1948, *AJ*, 107, 6
- Taylor, M., & Díaz, A. I. 2007, *PASP*, 374, 104
- Taylor, M., Díaz, A. I., Jodar-Sanchez, L. A., & Villanueva-Mico, R. J. 2008, *Adv. Studies Theor. Phys.*, 2, 979